


ILUVIEN®
PATIENT EDUCATIONAL MATERIALS



IMPORTANT SAFETY INFORMATION

You can access an audio version of this information by scanning the QR code in the top right-hand corner of this leaflet with a smartphone.

Alternatively you can request a large print version from your eye doctor or from Alimera via the contact details below.

IT IS IMPORTANT TO BE AWARE OF POTENTIAL CHANGES IN YOUR EYE AFTER THE ILUVIEN INJECTION

In rare cases, the injection can cause an infection in your eye or changes to the retina.

SEEK URGENT MEDICAL HELP IF YOU EXPERIENCE ANY OF THE FOLLOWING SIGNS OR SYMPTOMS:

- Hazy or decreased vision
- Eye pain
- Worsening redness of your eye
- A feeling of spots in front of your eye
- Increased sensitivity to light after your injection
- Flashing lights
- Partially blocked vision
- Distorted vision or other visual disturbance
- Swelling of the surface of the eye (corneal swelling)
- Change in the appearance of your eye at the front.

Injection with ILUVIEN may also cause the following:

- Increase in pressure inside your eye
- Clouding of your eye's natural lens (cataract)

Therefore, you must attend the visits at the clinic so that you can be monitored by your doctor regularly.

Please read the ILUVIEN Patient Information Leaflet for complete safety information.

This can be obtained from www.medicines.ie

If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet.

Doctor

Tel

You can also report any side effects directly to:

HPRA Pharmacovigilance

Website: www.hpra.ie

Side effects can also be reported to:

Alimera Sciences Europe Limited

Email: pvalimerasciences@alimerasciences.com